

Research Article

ON PRE- $T_{\frac{3}{4}}$ SPACES

Khadiga S. Abdeen and *Khadiga A. Arwini

Mathematics Department, Faculty of Science, Tripoli University, Tripoli-Libya.

Received 27th March 2022; Accepted 29th April 2022; Published online 31th May 2022

ABSTRACT

Since every topological space is pre- $T_{\frac{1}{2}}$ space, we aim to define a space that is weaker than pre- $T_{\frac{1}{2}}$ space; namely pre- $T_{\frac{3}{4}}$ space, when we use the notions of regular open sets and preopen sets. We prove that a pre- $T_{\frac{3}{4}}$ space is weaker than $T_{\frac{3}{4}}$ -space, regular space and pre- R_0 space, additionally, we investigate the topological properties of pre- $T_{\frac{3}{4}}$ space, as the hereditary property and their images by some particular functions; moreover we discuss the behavior of pre- $T_{\frac{3}{4}}$ axiom in some special spaces as; submaximal spaces, regular spaces, extremely disconnected spaces and hyperconnected spaces.

Keywords: Topological space and generalizations, generalized continuity, separation axioms, subspaces, submaximal spaces, regular spaces, hyperconnected space, extremely disconnected space.

AMS Subject Classification (2000): 54A05, 54C08, 54D70, 54B05, 54D10, 54D65.

INTRODUCTION

In 1937 Stone [1], introduced regular open sets using the closure and interior operators, and he showed that regular open set is stronger than open set, then he applied the family of regular open sets in the semi regularization space and in a generalization for algebraic openings and closings in a complete lattice. More details can be found in [2-8]. Corson and Michael in 1964 [9] defined the concepts of preopen sets by the name "locally dense" sets, as a generalizations of open sets, this class of sets plays a significant role in general topology. See more in [10-15]. In the literature many studies have been made on defining different separation axioms using some generalizations of open sets and closed sets, where the characterization of these spaces found to be useful in the study of computer science and digital topology. Levine, Maki, Dunham, Dontchev, Araki and Arenas, introduced several low separation axioms that lie between T_0 and T_1 -spaces, in particular, they used the concepts of λ -sets, generalized λ -sets, λ -sets, generalized closed sets and regular open sets to defined the axioms of $T_{\frac{1}{4}}$, $T_{\frac{1}{3}}$, $T_{\frac{1}{2}}$ and $T_{\frac{3}{4}}$ spaces [16-23]. The concept of generalized closed set was due to Levine in 1970 [16], when he used this notation to define a space called $T_{\frac{1}{2}}$ -space, and he showed that $T_{\frac{1}{2}}$ is strictly between the spaces T_1 and T_0 , in addition, he produced new separation axiom which lies between $T_{\frac{1}{2}}$ -space and T_1 -space, called $T_{\frac{3}{4}}$ -space. After that, the authors in [17] characterized $T_{\frac{1}{2}}$ -spaces as those spaces where every subset is λ -closed. Dunham [18] showed that a space is $T_{\frac{1}{2}}$ if and only if each singletons is open or closed.

Pre-separation axioms consist the spaces; pre- T_0 , pre- T_1 , pre- T_2 ,

pre-regular and pre-normal spaces and they were defined by many researchers as Kar and Bhattacharye [24], by replacing the notion of open sets with preopen sets in the classical definitions. In 1995 [25], Dontchev defined pre- $T_{\frac{1}{2}}$ space as a space in which every singleton is either preopen or preclosed, and he used the property that any singleton is either preopen or nowhere dense set which due to Jankovic and Reilly [26], to prove that any topological space is pre- $T_{\frac{1}{2}}$ space. Since any topological space is pre- $T_{\frac{1}{2}}$, we aim to define a space that is weaker than pre- $T_{\frac{1}{2}}$ space and $T_{\frac{3}{4}}$ -space; called pre- $T_{\frac{3}{4}}$ space, when we use the concepts of regular open sets and preopen sets. In this paper, we study the topological properties of this space, as the hereditary property and their images by some particular functions. Moreover, we characterized this space by the new notion of $pg\delta$ -closed set. Finally, we investigate the behavior of pre- $T_{\frac{3}{4}}$ space in some special spaces as; regular spaces, submaximal spaces, partition spaces, hyperconnected spaces and extremely disconnected spaces. We divided this article into six sections, as follows: in section one we present the introduction, then we review some preliminaries in section two, regarding regular open sets, preopen sets and some types of separation axioms that we need in our study. In section three, we define pre- $T_{\frac{3}{4}}$ space using the notion of regular open sets and preclosed sets, and then we study their relation with the classical separation axioms, and investigate the subspace behavior and their images by some particular maps. Section four contains a new concepts of $pg\delta$ -closed set as a generalization of closed set, and then we used this concept to study pre- $T_{\frac{3}{4}}$ space. In section five, we investigate the characterization of pre- $T_{\frac{3}{4}}$ spaces and establish their implication with some known spaces. Finally, we outline our results in the conclusion.

*Corresponding Author: Khadiga A. Arwini,
Mathematics Department, Faculty of Science, Tripoli University, Tripoli-Libya.

PRELIMINARIES

In this section, we recall some details that we need in the sequel regarding: regular open sets, preopen sets, some low separation axioms and pre-separation axioms. Throughout this paper (X, τ) or simply X denotes topological space, and we shall denote the closure and the interior of a set A with respect to the topological space (X, τ) as \bar{A} and A° ; respectively, and the complement of the set A in X , the difference of A and B , and the power set of X are denoted by A^c , A/B and $P(X)$; respectively.

Regular Open Sets and Preopen Sets

Definition 2.1.1. [8,11] A subset M of a space (X, τ) is called:

1. regular open (briefly r -open) if $W = \overline{W^\circ}$, while the set is called δ -open set if it is the union of r -open sets.
2. preopen if $W \subseteq \overline{W^\circ}$. The complement of preopen set is called preclosed.

The family of all regular open sets in X is denoted by $RO(X, \tau)$, while the family of all preopen sets and preclosed sets in X are denoted by $PO(X, \tau)$ and $PC(X, \tau)$, respectively.

Corollary 2.1.2. [8,11] In a space X , we have:

- every r -open set is δ -open set.
- every δ -open set is open.
- every open set is preopen.
- every denes set is preopen.

$$r\text{-open set} \implies \delta\text{-open set} \implies \text{open set} \implies \text{preopen set}$$

Corollary 2.1.3. [11] A subset B of a space X is preclosed iff $\overline{B^\circ} \subseteq B$.

Corollary 2.1.4. [8,11] R -closed sets, δ -closed sets, closed sets and preclosed sets are weakly ordered as:

$$r\text{-closed set} \implies \delta\text{-closed set} \implies \text{closed set} \implies \text{preclosed set}$$

Definition 2.1.5. [8,11] Let B be a subset of a topological space (X, τ) , then:

1. the pre-closure of B defined as the intersection of all preclosed sets containing B , and denoted by \overline{B}^p .
2. the δ -closure of B defined as the intersection of all δ -closed sets containing B , and denoted by \overline{B}^δ .

Theorem 2.1.6. [8,11] Let X be a space and $B \subseteq X$, then:

- $B \subseteq \overline{B}^p \subseteq \bar{B} \subseteq \overline{B}^\delta$.
- B is preclosed set if and only if $\overline{B}^p = B$.
- B is δ -closed set if and only if $\overline{B}^\delta = B$.
- \overline{B}^p is preclosed set, while \overline{B}^δ is δ -closed set.

Theorem 2.1.7. [8,11] If (X, τ) is a topological space, and $W \subseteq Y \subseteq X$; where:

- Y is an open subspace of X , then W is r -open in Y iff W is r -open in X .
- Y is an preopen subspace of X , then W is preopen in Y iff W is preopen in X .

Definition 2.1.8. [4,11] A map $F: (X, \tau) \rightarrow (Y, \sigma)$ is called:

1. Strongly regular open map if the image of each r -open subset of X is r -open in Y .
2. preclosed map if the image of each preclosed subset of X is preclosed in Y .

Separation Axioms

Definition 2.2.1. [27,16] A topological space (X, τ) is said to be:

1. T_1 -space if whenever x, y are distinct points in X , there exist open sets U and V such that U containing x but not y , and V containing y but not x .
2. $T_{\frac{1}{2}}$ -space if $\{x\}$ is open or closed for any point x in X .
3. $T_{\frac{3}{4}}$ -space if $\{x\}$ is r -open or closed for any point x in X .

Theorem 2.2.2. [27] A topological space X is T_1 -space if any singleton $\{x\}$ is closed in X .

Corollary 2.2.3. [16]

- Every T_1 -space is $T_{\frac{3}{4}}$.
- Every $T_{\frac{3}{4}}$ -space is $T_{\frac{1}{2}}$.

Definition 2.2.4. [24, 25] A topological space (X, τ) is said to be:

1. pre- T_1 space if whenever x, y are distinct points in X there exist preopen sets A and B such that A containing x but not y , and B containing y but not x .
2. pre- $T_{\frac{1}{2}}$ space if $\{x\}$ is preopen or preclosed for any point x in X .
3. pre- T_0 space if whenever x, y are distinct points in X there exists a preopen set that containing one and not the other.

Theorem 2.2.5. [24] A space X is pre- T_1 iff for any point $x \in X$, the singleton set $\{x\}$ is preclosed.

Corollary 2.2.6. [24] Every T_1 -space is pre- T_1 , where $(i=0, \frac{1}{2}, 1)$.

Definition 2.2.7. [24] A space X is said to be pre- R_0 iff for each preopen set G and $x \in G$ implies $\overline{\{x\}}^p \subseteq G$.

Corollary 2.2.8. [24] In a space X , we have:

- Any pre- T_1 space is pre- R_0 .
- In pre- R_0 space, X is pre- T_1 iff X is pre- T_0 .

PRE- $T_{\frac{3}{4}}$ SPACES

In this section, we define a new space that is weaker than both $T_{\frac{3}{4}}$ -space and pre- T_1 space; called pre- $T_{\frac{3}{4}}$ space, then we study the topological properties of this new axiom as their implication with the classical separation axioms, additionally, we discuss their subspaces behavior and their images by some particular maps.

Definition 3.1. A topological space (X, τ) is called pre- $T_{\frac{3}{4}}$ space if every singleton is regular open or preclosed.

Theorem 3.2. Every pre- T_1 space is pre- $T_{\frac{3}{4}}$.

Proof. From theorem (2.2.5), every singleton is preclosed set in pre- T_1 space, it follows that the space is pre- $T_{\frac{3}{4}}$.

$$\text{Pre-}T_1 \text{ Space} \implies \text{Pre-}T_{\frac{3}{4}} \text{ Space}$$

Theorem 3.3. Every $T_{\frac{3}{4}}$ -space is $p-T_{\frac{3}{4}}$.

Proof. From definition (2.2.1) and corollary (2.1.4) any singleton $\{x\}$ is r -open or preclosed in $T_{\frac{3}{4}}$ -space, it follows that the space is $p-T_{\frac{3}{4}}$.

$$T_{\frac{3}{4}}\text{-Space} \implies \text{Pre-}T_{\frac{3}{4}}\text{ Space}$$

Examples 3.4.

1. If $X = \{a,b,c\}$ with $\tau = \{\emptyset, X, \{a,b\}\}$, then $RO(X, \tau) = \{\emptyset, X\}$ and $PO(X, \tau) = P(X)/\{\{c\}\}$. Therefore, the space X is $p-T_{\frac{3}{4}}$ but not $T_{\frac{3}{4}}$ -space since the singleton $\{a\}$ is not r -open nor closed.
2. Let $X = \mathbb{R}$ with $\tau = \{A \subseteq \mathbb{R}: 0 \notin A\} \cup \{\emptyset\}$, then $PO(X, \tau) = RO(X, \tau) = \tau$, so any singleton $\{x\}$ where $x \neq 0$ is r -open, and $\{0\}$ is preclosed. Thus X is $p-T_{\frac{3}{4}}$ space.
3. Let $X = \{a,b,c\}$ with $\tau = \{\emptyset, X, \{a\}\}$, the $RO(X, \tau) = \{X, \emptyset\}$ and $PO(X, \tau) = \{\emptyset, X, \{a\}, \{a,b\}, \{a,c\}\}$, so X is not $p-T_{\frac{3}{4}}$, since the singleton $\{a\}$ is not r -open nor preclosed.

Lemma 3.5. In a space X , any singleton $\{x\}$ is r -open iff $\{x\}$ is δ -open.

Proof. Suppose $\{x\}$ is δ -open, then $\{x\} = \bigcup_{\alpha} W_{\alpha}$; where W_{α} is r -open set, hence there is α such that $W_{\alpha} = \{x\}$; so $\{x\}$ is r -open. Conversely, direct since any r -open set is δ -open.

$$\{x\} \text{ is } r\text{-open} \implies \{x\} \text{ is } \delta\text{-open}$$

Corollary 3.6. A space X is $p-T_{\frac{3}{4}}$ iff any singleton $\{x\}$ is δ -open or preclosed.

Proof. Direct from the previous lemma.

Lemma 3.7. If (X, τ) is a topological space, then any singleton $\{x\}$ is open or preclosed.

Proof. Suppose $\{x\}$ is not open in X , then $\{x\}^c$ is not closed, i.e. $\overline{\{x\}^c} \neq \{x\}^c$ hence $\overline{\{x\}^c} = X$, so $\overline{\{x\}^c}^{\circ} = \overline{X}^{\circ} = X$, therefore $\{x\}^c$ is preopen set, hence $\{x\}$ is preclosed.

Theorem 3.8. If (X, τ) is a $p-T_{\frac{3}{4}}$ space and $Y \subseteq X$ is an open subspace of X , then Y is $p-T_{\frac{3}{4}}$.

Proof. Let $y \in Y$, since X is $p-T_{\frac{3}{4}}$, then $\{y\}$ is r -open or preclosed set in X . If $\{y\}$ is r -open in X , then from theorem (2.1.7) we obtain $\{y\}$ is r -open in Y . If $\{y\}$ is preclosed in X , then from corollary (2.1.3) we have $\overline{\{y\}}^{\circ} \subseteq \{y\}$: **Case one** when $\{y\}$ is not open in X , then $\{y\}$ is not open in Y (since Y is open subspace), then from lemma (3.7) we obtain $\{y\}$ is preclosed in Y . **Case two** when $\{y\}$ is open in X , and from $\overline{\{y\}}^{\circ} \subseteq \{y\}$ we obtain $\overline{\{y\}} \subseteq \{y\}$, hence $\overline{\{y\}} = \{y\}$, i.e. $\{y\}$ is open and closed, so $\{y\}$ is clopen in X , then $\{y\}$ is also clopen in Y (since $\{y\} = \{y\} \cap Y$), hence $\{y\}$ is preclosed in Y . Thus complete the prove.

Example 3.9. Subspace of $p-T_{\frac{3}{4}}$ space need not be $p-T_{\frac{3}{4}}$ in general, for example if $X = \{a,b,c\}$ with $\tau = \{\emptyset, X, \{a,b\}\}$, then the space X is $p-T_{\frac{3}{4}}$ while the subspace $Y = \{b,c\}$ where $\tau_Y = \{\emptyset, Y, \{b\}\}$ is not $p-T_{\frac{3}{4}}$, since the singleton $\{b\}$ is not r -open nor preclosed in Y .

Theorem 3.10. The image of a $p-T_{\frac{3}{4}}$ space by preclosed and strongly regular open map is $p-T_{\frac{3}{4}}$ space.

Proof. Let $F: (X, \tau) \rightarrow (Y, \sigma)$ be an onto map from a $p-T_{\frac{3}{4}}$ space X , and suppose $y \in Y$, then there exists a point $x \in X$ such that $F(x) = y$, since X is $p-T_{\frac{3}{4}}$ we have $\{x\}$ is r -open or preclosed in X , since the map F is preclosed and strongly regular open, we have $F(\{x\}) = \{y\}$ is r -open or preclosed in Y .

Example 3.11. Strongly regular open image of $p-T_{\frac{3}{4}}$ space need not be $p-T_{\frac{3}{4}}$ space, for example: Let $X = Y = \mathbb{R}$ and let τ_1 be the trivial topology on \mathbb{R} , and $\tau_2 = \{\emptyset, \mathbb{R}, \{1\}\}$. Then $PO(\mathbb{R}, \tau_1) = P(\mathbb{R})$ and $RO(\mathbb{R}, \tau_1) = \{\emptyset, \mathbb{R}\}$, while $PO(\mathbb{R}, \tau_2) = \{A \subseteq \mathbb{R}: 1 \in A\}$ and $RO(\mathbb{R}, \tau_2) = \{\emptyset, \mathbb{R}\}$. Then the identity map from (\mathbb{R}, τ_1) onto the space (\mathbb{R}, τ_2) is strongly regular open map, however the space (\mathbb{R}, τ_1) is $p-T_{\frac{3}{4}}$, since, while (\mathbb{R}, τ_2) is not $p-T_{\frac{3}{4}}$ space, since $\{1\}$ is not r -open nor preclosed.

PRE-GENERALIZED δ -CLOSED SETS AND $T_{\frac{3}{4}}$ -SPACES

In this section, we introduce a new generalization of δ -closed sets using the notions of preopen sets; namely pre-generalized δ -closed sets. We investigate their properties and characterized the space of $p-T_{\frac{3}{4}}$ using this class of sets.

Definition 4.1. A subset B of a topological space (X, τ) is called pre-generalized δ -closed (pg δ -closed) if $\overline{B}^{\delta} \subseteq N$ whenever $B \subseteq N$ and N is preopen set.

Corollary 4.2. Any δ -closed set is pg δ -closed, but not conversely.

Proof. Suppose B is δ -closed set and $B \subseteq N$, where N is preopen, then $\overline{B}^{\delta} \subseteq B \subseteq N$, hence B is δ g-preclosed.

$$\delta\text{-closed set} \implies \text{pg}\delta\text{-closed set}$$

Example 4.3. Let $X = \{a,b,c\}$ with $\tau = \{\emptyset, X, \{a\}, \{b\}, \{a,b\}, \{b,c\}\}$, then $A = \{c\}$ is pg δ -closed but it is not δ -closed.

Examples 4.4. No general relation between closed set and pg δ -closed set, for example:

1. If $X = \mathbb{R}$ with $\tau = \{\mathbb{R}, \mathbb{Q}, \emptyset\}$, then $RO(X, \tau) = \{\mathbb{R}, \emptyset\}$ while $PO(X, \tau) = \{A \subseteq \mathbb{R}: A \cap \mathbb{Q} \neq \emptyset\} \cup \{\emptyset\}$, so the set \mathbb{K} is closed but not pg δ -closed, since $\mathbb{K} \subseteq \mathbb{K} \cup \{2\}$ and $\mathbb{K} \cup \{2\}$ is preopen set but $\overline{\mathbb{K}}^{\delta} = \mathbb{R} \not\subseteq \mathbb{K} \cup \{2\}$.
2. In the usual topology on \mathbb{R} the interval $[0,1)$ is pg δ -closed set but not closed, since $\overline{[0,1)}^{\delta} = [0,1]$, and the smallest preopen set that contains $[0,1)$ is the closed interval $[0,1]$.

Theorem 4.5. A space X is $p-T_{\frac{3}{4}}$ iff any pg δ -closed set is δ -closed.

Proof. Suppose A is pg δ -closed set but not δ -closed, i.e. $\overline{A}^{\delta} \neq A$, hence there exists a point $x \in \overline{A}^{\delta}$ but $x \notin A$, then any δ -open set contains x intersects A , but $\{x\} \cap A = \emptyset$, hence $\{x\}$ is not δ -open, so it is not r -open, and since X is $p-T_{\frac{3}{4}}$, we have $\{x\}$ is preclosed.

Then $A \subseteq \{x\}^c$, where $\{x\}^c$ is preopen, so $\overline{A}^{\delta} \subseteq \{x\}^c$, i.e. $x \notin \overline{A}^{\delta}$ which contradict the assumption. Conversely, suppose $\{x\}$ is not preclosed set, then $\{x\}^c$ is not preopen, so $\{x\}^c$ is pg δ -closed set

since the only preopen set contains $\{x\}^c$ is X , hence $\{x\}^c$ is δ -closed, i.e. $\{x\}$ is δ -open, therefore X is $\text{pre-}T_{\frac{3}{4}}$ -constant.

PRE- $T_{\frac{3}{4}}$ SPACES AND SOME RELATED SPACES

In the present section, we illustrate the implication between $\text{pre-}T_{\frac{3}{4}}$ space with some known spaces as; submaximal space, regular space, partition space, hyperconnected space and extremely disconnected space.

In Submaximal Spaces

Definition 5.1.1. [11] A space X is called submaximal if any dense set in X is open.

Theorem 5.1.2. [11] space X is submaximal iff any preopen is open.

Corollary 5.1.3. In submaximal space X , these conditions are equivalent:

- X is $\text{pre-}T_{\frac{3}{4}}$.
- X is $T_{\frac{3}{4}}$.

Proof. Direct since any preopen set is open.

$$\text{Pre-}T_{\frac{3}{4}} \text{ Space} \xleftrightarrow{\text{Submaximal}} T_{\frac{3}{4}}\text{-Space}$$

Corollary 5.1.4. In submaximal space X , we have:

1. X is $\text{pre-}T_0$ iff X is T_0 .
2. X is $\text{pre-}T_1$ iff X is T_1 .
3. X is $\text{pre-}T_2$ iff X is T_2 .
4. X is $\text{pre-}T_{\frac{1}{2}}$ iff X is $T_{\frac{1}{2}}$.

Corollary 4.1.5. Any submaximal space is $T_{\frac{1}{2}}$.

Proof. Since any space is $\text{pre-}T_{\frac{1}{2}}$ and in submaximal space, the axioms $\text{pre-}T_{\frac{1}{2}}$ and $T_{\frac{1}{2}}$ are coincid, thus complete the prove.

$$\text{Submaximal Space} \implies T_{\frac{1}{2}}\text{-Space}$$

In Regular Spaces

Theorem 5.2.1. [6] In regular space X , any open set can be expressed as a union of r -open sets, equivalently; a subset A of a regular space X is open set if and only if A is δ -open

Corollary 5.2.2. In regular space X , the singleton $\{x\}$ is r -open iff $\{x\}$ is open.

Proof. From corollary (2.1.2) any r -open set is open. Conversely, any open set in regular space is δ -open, and from lemma (3.5) any δ -open singleton is r -open set, that complete the prove.

Theorem 5.2.3. Any regular space X is $\text{pre-}T_{\frac{3}{4}}$.

Proof. From lemma (3.7) any singleton $\{x\}$ is open or preclosed, and in regular space any singleton open set is r -open, so we have $\{x\}$ is r -open or preclosed, hence X is $\text{pre-}T_{\frac{3}{4}}$.

$$\text{Regular Space} \implies \text{Pre-}T_{\frac{3}{4}} \text{ Space}$$

Example 5.2.4. $\text{Pre-}T_{\frac{3}{4}}$ space need not be regular, for example: Let $X = \{a, b, c\}$ with $\tau = \{\emptyset, X, \{a, b\}\}$, then $\text{RO}(X, \tau) = \{\emptyset, X, \{a, b\}, \{a\}, \{b\}, \{a, c\}, \{b, c\}\}$ and

$\text{PC}(X, \tau) = \{\emptyset, X, \{a\}, \{b\}, \{c\}, \{b, c\}, \{a, c\}\}$. Therefore the space X is $\text{p-}T_{\frac{3}{4}}$ but not regular.

Corollary 5.2.5 In regular-submaximal space X , these statements are equivalent:

1. X is $\text{pre-}T_{\frac{3}{4}}$.
2. X is $T_{\frac{3}{4}}$.
3. X is $T_{\frac{1}{2}}$.

Proof.

1. $\implies 2$) Direct since in submaximal space any preclosed set is closed.
2. $\implies 3$) Direct.
3. $\implies 1$) Direct since in regular space any singleton open set is r -open.

$$\text{Pre-}T_{\frac{3}{4}} \text{ Space} \xleftrightarrow{\text{Regular submaximal}} T_{\frac{3}{4}}\text{-Space} \xleftrightarrow{\text{Regular submaximal}} T_{\frac{1}{2}}\text{-Space}$$

Theorem 5.2.6. Any regular submaximal space X is $T_{\frac{3}{4}}$.

Proof. Since any regular space X is $\text{pre-}T_{\frac{3}{4}}$, and any $\text{pre-}T_{\frac{3}{4}}$ submaximal space is $T_{\frac{3}{4}}$, thus complete the prove.

$$\text{Regular submaximal Space} \implies T_{\frac{3}{4}}\text{-Space}$$

In Partition Spaces

Definition 5.3.1. [14] A space (X, τ) is called partition if any open set is closed.

Remark 5.3.2. In partition space (X, τ) , we have $\text{RO}(X, \tau) = \tau$ and $\text{PO}(X, \tau) = \text{P}(X)$.

Proof. Suppose V is a non-empty open subset of X , then $\overline{V}^o = V^o = V$, so $\text{RO}(X, \tau) = \tau$. Moreover, $N \subseteq \overline{N}$ for any subset N of X , since in partition space any closed set is open, we get $\overline{N}^o = \overline{N}$, hence $N \subseteq \overline{N}^o$, i.e. $\text{PO}(X, \tau) = \text{P}(X)$.

Corollary 5.3.3. Any partition space (X, τ) is $\text{p-}T_{\frac{3}{4}}$ (and $\text{pre-}T_1$).

Proof. Any singleton in partition space is preclosed, since $\text{PO}(X, \tau) = \text{P}(X)$.

$$\text{Partition Space} \implies \text{Pre-}T_{\frac{3}{4}} \text{ Space}$$

Corollary 5.3.4. Any $T_{\frac{1}{2}}$ partition space is $T_{\frac{3}{4}}$.

Proof. Since any singleton in $T_{\frac{1}{2}}$ -space is open or closed, and in partition space any open is r -open, then X is $T_{\frac{3}{4}}$.

$$T_{\frac{1}{2}}\text{-Space} \xleftrightarrow{\text{Partition}} T_{\frac{3}{4}}\text{-Space}$$

In Hyperconnected Spaces

Definition 5.4.1. [14] A space X is called hyperconnected if any open set is dense.

Lemma 5.4.2. A space X is called hyperconnected space (X, τ) , we have $\text{RO}(X, \tau) = \{\emptyset, X\}$.

Proof. Suppose V is a non-empty open set, then V is dense in X , so $\overline{V}^o = X^o = X$, hence V is not r -open set.

Theorem 5.4.3. Any pre- $T_{\frac{3}{4}}$ hyperconnected space X is pre- T_1 .

Proof. Suppose X is pre- $T_{\frac{3}{4}}$, then any singleton $\{x\}$ is r -open or preclosed, but from the previous lemma $\{x\}$ is not r -open, hence $\{x\}$ is preclosed, thus X is pre- T_1 .

Corollary 5.4.3. In hyperconnected space X , we have: X is pre- $T_{\frac{3}{4}}$ iff X is pre- T_1 .

$$\text{Pre-}T_{\frac{3}{4}} \text{ Space} \xleftrightarrow{\text{Hyperconnected}} \text{Pre-}T_1 \text{ Space}$$

In Extremely Disconnected Space

Definition 5.5.1. [14] A topological space (X, τ) is called extremely discometed (namely e.d) if the closure of any open set is open, i.e. if V is open in X , then \bar{V} is also open.

Corollary 5.5.2. [14] A space (X, τ) is extremely discometed, then:

1. any r -open set is clopen.
2. any r -closed set is clopen .
3. $RO(X, \tau) = RC(X, \tau) = \tau \cap F$, where F is the collection of all closed subsets in X .

Theorem 5.5.3. Any pre- $T_{\frac{3}{4}}$ extremely discometed space is pre- R_0 .

Proof. Suppose N is preopen set in X and $x \in N$, since X is pre- $T_{\frac{3}{4}}$ then $\{x\}$ is r -open or preclosed. **Case 1:** If $x \in N$ and $\{x\}$ is r -open, then since X is extremely discometed the singleton $\{x\}$ is clopen, so $\{x\}$ is preclosed, i.e. $\overline{\{x\}}^p = \{x\} \subseteq N$, hence X is pre- R_0 . **Case 2:** If $x \in N$ and $\{x\}$ is preclosed, then $\overline{\{x\}}^p = \{x\} \subseteq N$, hence X is pre- R_0 .

$$\text{Pre-}T_{\frac{3}{4}} \text{ Space} \xrightarrow{\text{Extremely disconnected}} \text{Pre-}R_0 \text{ Space}$$

Theorem 5.5.3. Any pre- R_0 space is pre- $T_{\frac{3}{4}}$.

Proof. If X is a space, then X is pre- $T_{\frac{1}{2}}$, so it is pre- T_0 space, and from corollary (2.2.8) any pre- T_0 pre- R_0 is pre- T_1 .

Corollary 5.5.4. Any pre- R_0 space is pre- T_1 .

$$\text{Pre-}R_0 \text{ Space} \implies \text{Pre-}T_1 \text{ Space} \implies \text{Pre-}T_{\frac{3}{4}} \text{ Space}$$

Remark 5.5.5. In extremely disconnected space X , we have: X is pre- $T_{\frac{3}{4}}$ iff X is pre- R_0 .

$$\text{Pre-}T_{\frac{3}{4}} \text{ Space} \xleftrightarrow{\text{Extremely disconnected}} \text{Pre-}R_0 \text{ Space}$$

CONCLUSION

Using the concept of regular open sets and preclosed sets we introduce a new space namely pre- $T_{\frac{3}{4}}$ space, this space is weaker than both pre- T_1 space and $T_{\frac{3}{4}}$ -space. In this paper, we study the topological properties of this space, as the hereditary property and their images by some particular functions, moreover, we characterized this space by the new notion of $pg\bar{\delta}$ -closed sets. Finally, we investigate the behavior of pre- $T_{\frac{3}{4}}$ space in some special spaces as; regular spaces, submaximal spaces, partition spaces, hyperconnected spaces and extremely disconnected spaces. Outline some of our results:

- A space X is pre- $T_{\frac{3}{4}}$ iff any $pg\bar{\delta}$ -closed set is $\bar{\delta}$ -closed set.

- Pre- $T_{\frac{3}{4}}$ space is weaker than: $T_{\frac{3}{4}}$ -space, partition space, regular space, pre- T_1 space and pre- R_0 space.
- In hyperconnected space; the spaces pre- $T_{\frac{3}{4}}$ space and pre- T_1 space are coincide.
- In extremely disconnected space; the spaces pre- $T_{\frac{3}{4}}$ space and pre- R_0 space are coincide.
- In submaximal space; the spaces pre- $T_{\frac{3}{4}}$ space and $T_{\frac{3}{4}}$ -space are coincide.
- Any regular submaximal space is $T_{\frac{3}{4}}$ -space.

REFERENCES

1. M. H. Stone, Applications of The Theory of Boolean Rings to General Topology. Trans. Am. Math. Soc., 41(1937) 375-481.
2. C. Ronse, Regular Open and Closed Sets. Philips Research Laboratory Brussels, Ar. E Van Baclaera, 2 B (1990) 11-70.
3. Charles Dorsett, Regularly Open Sets and R-Topological Properties. Nat. Acad. Sci-Letters, 10 (1987) 17-21.
4. H. K. Abdullah and F. K. Radhy, Strongly Regular Proper Mappings. Journal of Al- Qadisiyah for Computer Science and Mathematics, 3 (1) (2011).
5. Zanyar A. Ameen, Baravan A. Asaad and Ramadhan A. Muhammed, On Super Classes of Open Sets in Topological Spaces. International Journal of Applied Mathematics, 32 (2) (2019) 259-277.
6. Amel E. Kornas and K. A. Arwini, R-Countability Axioms. An International Scientific Journal, 149 (2020) 92-109.
7. Khadiga A. Arwini and Entisar O. Leghah, λ -Generalizations and g -Generalizations. Tarbawe Journal, Elmergib, 19 (2021) 245-255.
8. Mabrouka M. Almarghani and Khadiga A. Arwini, Generalizations of Regular Closed Sets. An international Scientific Journal, 152 (2021) 55-68.
9. H. H. Corson & E. Michael, Metrizable of Certain Countable Unions. Illinois J. Math., 8 (2) (1964) 351-360.
10. Frank Siwec, On Defining A Space By A Weak Base. Pacific Journal of Mathematics, 52 (1) (1974) 233-245.
11. J. Dontchev, Survey On Preopen Sets. The Proceedings of The Yatsushiro Topological Conference, (1998) 11-18.
12. Young Bae Jun, Seong Woo Jeong, Hyeon Jeong Lee and Joon Woo Lee, Application of Preopen Sets. Applied General Topology, Universidad Politecnica De Valencia, 9 (2) (2008) 213-228.
13. Raad Aziz Hussain Al-Abdulla, Preopen Sets in Topological Spaces. Al-Qadisiyah Journal of Pure Science, 12 (4) (2007) 203-213.
14. Nagah A. Elbhillil and Khadiga A. Arwini, Countability Axioms Via Preopen Sets. An International Scientific journal, 152 (2021) 111-125.
15. V. V. S. Ramachandram, A Note On Preopen Sets. Research Journal of Pure Algebra, 1 (5) (2011) 132-134.
16. N. Levine, Generalized Closed Sets in Topology. Rend. Circ. Mat. Palermo, 19 (2) (1970) 89-96.
17. H. Maki, J. Umehara and K. Yamamura, Characterizations of $T_{\frac{1}{2}}$ -Spaces Using Generalized V-Sets. Indian J. Pure Appl. Math., 19 (7) (1988) 634-640.
18. W. Dunham, $T_{\frac{1}{2}}$ -Space. Kyungpook Math J., (17) (1977) 161-169.
19. N. Araki, M. Ganster, H. Maki and A. Nishi, Properties of $T_{\frac{1}{4}}$ Spaces. Questions and Answers in General Topology, 24 (2) (2006) 143-148.

20. Francisco G. Arenas, Julian Dontchev and Maximilian Ganster, On λ -Sets and the Dual of Generalized Continuity. Questions Answers in Gen Topology, 15 (1) (1997) 3–13.
21. F.G. Arenas, J. Dontchev and M.L. Puertas, Unification Approach to the Separation Axioms Between T_0 and Completely Hausdorff. Acta Math. Hungar, 86 (2000) 75-82.
22. Francisco G. Arenas, Julian Dontchev and Maria Luz Puertas, Idealization of Some Weak Separation Axioms. Acta Mathematica Hungarica, Source, 89 (2000) 47-53.
23. J. Dontchev and M. Ganster, On Minimal Door, Minimal Anti Compact and Minimal $T_{\frac{3}{4}}$ -spaces. Mathematical Proceedings of the Royal Irish Academy, 98 A (2) (1998) 209-215.
24. A. Kar and P. Bhattacharyye, Weak Separation Axioms in Terms of Preopen Sets. Bull. Cal. Math. Soc., 82 (1990) 415-422.
25. Julian Dontchev, On Door Spaces. Indian J. Pure Appl. Math., 26 (9) (1995) 873-881.
26. D. Jankovic and I. Reilly, On Semi-Separation Properties. Indian J. Pure Appl. Math., 16 (1985) 957-964.
27. N. Bourbaki, General Topology. Addison Wesley, Part 1, Reading, Massachusetts, (1966).
